

Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/SE04/001883

International filing date: 16 December 2004 (16.12.2004)

Document type: Certified copy of priority document

Document details: Country/Office: SE
Number: 0400191-3
Filing date: 30 January 2004 (30.01.2004)

Date of receipt at the International Bureau: 03 January 2005 (03.01.2005)

Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse

Ink. t. Patent- och reg.verket

+46 31630263

-G1- 3 0

Huvudfaxen Kassan

7

5 It is clear that even though it is well established in the scientific community that periodontal disease is a multi-factorial disease no one has yet developed a test to analyse both bacterial and host derived products in a sample.

10 The test kit may also comprise a third detection assay comprising at least a third affinity ligand having a binding site for binding of a third substance originating from bacteria or from the immune or inflammatory system of the patient as defined above, preferably from bacteria, and may in some embodiments even comprise further detection assays for binding of additional substances.

15 Preferably, the first affinity ligand is an antibody exhibiting selective binding of said first substance, and the second affinity ligand is an antibody exhibiting selective binding of said second substance.

20 The use of antibodies is advantageous over other methods (e.g. enzymatic methods) since antibodies, once developed and tested for cross-reactivity, are highly specific for their target and are able to detect chemical substances other than enzymes, e.g. toxins and cytokines. Furthermore, methods for development of new antibodies against new antigens are well known to those skilled in the art.

25 Preferably said detection assays in the test kit according to the present invention comprises immunochromatographic assays.

30 Among the advantages of using immunochromatographic assays/methods is that they are easily produced and used, have a long shelf-life, yields a quick answer and can be designed to be very specific for the substances intended to be detected.

35 Said test kit further preferably comprises a support provided with a sample reservoir for receiving a sample, wherein said first and second detection assays are arranged on said support in contact with said sample reser-

Ink. t. Patent- och reg.verket +46 31630263

-01- 3 0

Huvudfaxen Kassan

11

oides forsythus, or a toxin, more preferably the leukotoxin from *Actinobacillus actinomycetemcomitans*.

The presence of virulence products may trigger an inflammatory response and the immune system of the host, thus recruiting defence system cells, e.g. polymorphonuclear (PMN) leukocytes, to the site of infection.

The leukocytes cannot cope with the high amounts of bacteria and bacterial products at the infected site and enzymes from the leukocyte granulates are released into the periodontal pocket. These enzymes, originally aimed at killing the invading bacteria, are highly destructive to the surrounding tissues. Human neutrophil elastase has been shown to degrade many of the supportive tissues surrounding a tooth. Elastase is often found bound to its protease inhibitor (α -1 antitrypsin) in the periodontal pocket. However, the protease from *P. gingivalis* has been shown to degrade the protease inhibitors α -1 antitrypsin and α -2 macroglobulin in human serum, leaving elastase in its highly destructive form in the periodontal or peri-implant pocket.

As used herein, substances "originating from the immune or inflammatory system" refers to substances that originates from cells involved in the immune or inflammatory system. Such substances may be secreted from said cells, or may originate from lysis of such cells, for example to orchestrate the immune and inflammatory response or to remodel the tissue or killing the invading bacteria.

The second substance may be a leukocyte product, such as a natural serine protease, preferably human neutrophil elastase or a cytokine, such as an interleukin preferably chosen from among interleukin-1 β , interleukin-6 and interleukin-8, or an inflammatory mediator, preferably tumour necrosis factor- α or possibly prostaglandin E₂.

Most preferably, said second substance is human neutrophil elastase.

[illegible]

+46 31630263

Ink t. Patent- och reg.verket

-01-30

Huvudkassen Kassan

14

The invention also relates to a method for diagnosing periodontal diseases and/or predicting the risk for progress of said diseases, said method comprising analyzing a sample from the oral cavity of a patient for the presence of at least a first substance originating from bacteria and the presence of a second substance originating from the immune or inflammatory system of the patient.

In the method according to the present invention, sample from the oral cavity of a patient and the first and second substances are as defined above.

Also in some instances, the diagnostic method according to the present invention further comprises the step of detecting the presence of additional substances as defined above in said sample.

In preferred embodiments of the diagnosis method according to the present invention, said first method comprises using a first antibody exhibiting selective binding of said first substance and said second method comprises using a second antibody exhibiting selective binding of said second substance, wherein said antibodies are as defined above.

Most preferably at least one of said first and second methods comprises using an immunochromatographic method.

Other detection methods suitable for use with the methods according to the present invention may be based on one out of several immunological methods. Such methods include, but are not limited to, immunochromatographic methods, immunometric methods, immunoagglutination methods, fluoroimmunological methods, immunoluminescence methods, turbidimetric immunological methods, ELISA and nephelometric methods.

35 Preferred embodiments

Preferably a test kit according to the invention, as illustrated in figure 1, comprises two immunochroma-

Ink t. Patent- och reg.verket +46 31630263

-01-30

Huvudfaxen Kassan

15

tographic detection assays 1, 2 arranged on a support 3 equipped with a sample reservoir 4 for receiving a sample. The two detection assays 1, 2 are arranged, directly or via a removably arranged separating means 5, in contact with the sample reservoir 4.

The sample reservoir 4 is preferably formed in the support material. The support 3 can be made of several different material, such as plastic, paper, carton or combinations thereof, such as paper laminated with plastic. The detection assays 1, 2 are fixed on the support 3 in such a manner that the sample receiving areas of each assay is, directly or via a removably arranged separating means 5, in contact with the sample reservoir 4. Said separating means 5 may be a removable foil covering the sample receiving areas on the assays, a dam separating the reservoir from the assays, etc.

The kit may further comprise additional buffers for dilution and adaptation of said sample for said detection assays and at least one sampling device for obtaining the sample to be analysed. The type of sampling device suitable for use may depend on the type of sample to be analysed. For example gingival crevicular fluid is a viscous fluid and can easily be collected by a small brush, a dental floss, a paper point or a disposable pipette. Other sampling devices for obtaining a saliva or mouth rinse sample are known to those skilled in the art. Preferably the buffer is kept separately from the sample reservoir in a buffer reservoir, such as a separate flask, an additional reservoir formed on the support or by providing the buffer in a puncturable bag placed in the sample reservoir.

After a sample is obtained, it is preferably mixed with buffer to obtain the pH, ionic strength and viscosity suitable for the detection assays. After mixing, the sample is transferred to the sample reservoir, and is thus, optionally by removing the separating means,

'04 01/30 14:25 FAX +46 31630263

AWAPATIAAT

→ PRV KASSAN

020

Ink i Patent- och reg.verket

+46 31630263

-01 3 0

Huvudtaxen Kassan

18

(Nunc, Roskilde, Denmark) containing 100 µl 0,85% NaCl. The samples were frozen immediately at -20°C and within 6 hours at -80°C, and stored until assayed. For the microbial analysis 3 paper points were inserted consecutively into the periodontal pocket until resistance was met and left in place for 15 seconds. The points were pooled into a vial containing 10 glass beads, 3 mm in diameter, and 3,3 ml of VMGA III transport medium, aerobically prepared and stored. The samples were processed within 24 hours. Bacteria were grown on enriched Brucella agar plates and identified by appropriate methods.

For the study of appropriate cut-off values for a diagnostic test we studied elastase from human neutrophils and arg-gingipain from *P. gingivalis* using selective substrates.

The first set of the duplicate samples was used for the enzyme assays. All samples were thawed on ice and centrifuged for 3 minutes at 13,000 × g.

The enzyme substrate for determining arg-gingipain from *P. gingivalis* was N-benzoyl-L-arginine-p-nitroanilic (BAPNA) with a final concentration of 1 mM in the assay buffer containing 5 % DMSO. The assay buffer was 0.1 M Tris-HCl containing 5 mM CaCl₂, pH 7,5, with 50 mM glycyl-glycine (as reported earlier by the inventors in patent US 5,981,164 gly-gly stimulates BAPNA selectively in the presence of arg-gingipain) and 5 mM L-cysteine. Ten µL of the sample was preincubated for 15 minutes with 140 µL of the assay buffer in the well of a 96-well microtiter plate, precoated with bovine serum albumin, before 50 µL of the substrate was added. The plate was incubated at 37°C in a humid chamber, and the release of pNA was followed spectrophotometrically by OD₄₀₅ readings, using a microtiter plate reader, every few hours from 12 to 36 hours. One unit of activity was equal to the amount of enzyme which cleaved 1 nmol of the substrate during one hour of incubation.

+46 31630263

Ink t. Patent- och reg.verket

-C1- 3 0

Huvudtaxen Kassan

20

Results

| <u>Elastase</u> | | Attachment loss | Attachment gain or zero |
|---------------------------------|--|-----------------|-------------------------|
| Positive test (Elastase >20 ng) | | 3 | 2 |
| Negative test (Elastase ≤20 ng) | | 3 | 27 |

- 5 Elastase as a predictor of further attachment loss yielded a sensitivity of 50% and a specificity of 93%. A p-value of 0.0264* was calculated using Fisher's exact test.

| <u>Arg-gingipain</u> | | Attachment loss | Attachment gain or zero |
|---------------------------------------|--|-----------------|-------------------------|
| Positive test (arg-gingipain >0,27 U) | | 6 | 13 |
| Negative test (arg-gingipain ≤0,27 U) | | 0 | 16 |

10

Arg-gingipain as a predictor of further attachment loss yielded a sensitivity of 100% and a specificity of 55%. A p-value of 0.0216* was calculated using Fisher's exact test.

15

For elastase and arg-gingipain alone either the sensitivity or the specificity had to be sacrificed. This prompted us to investigate the combination of the two enzymes. We investigated new cut-off levels for the two enzymes to find that the detection limit for elastase should be 2 ng per site and 0,30 units for arg-gingipain.

20

+46 31630263

Ink. t. Patent- och reg.verket

2014-01-30

21

Huvudtaxen Kassan

| <u>Combination</u> | | Attachment loss | Attachment gain or zero |
|---|--|--------------------|----------------------------|
| Positive test (Elastase >2 ng <u>and</u> arg-gingipain >0,30 U) | | 5 | 3 |
| Negative test (Elastase ≤2 ng <u>or</u> arg-gingipain ≤0,30 U) | | 1 | 26 |

5 The combination of elastase and arg-gingipain as a
predictor of further attachment loss yielded a
sensitivity of 83% and a specificity of 90%. A p-value of
less than 0,001*** was calculated using Fisher's exact
test. This limited data shows that the combination of
elastase and arg-gingipain as a marker for periodontal
disease yields a statistically more significant test
10 than either of the enzymes alone.

+46 31630263

Ink. t. Patent och ren.verket

2001-01-30

22 Huvudstaden Kassar

References

- Armitage G.C. (2003) Position paper: Diagnosis of Periodontal Diseases. Journal of Periodontology 74, 1237-1247.
- 5 Eley B.M. and Cox S.W. (1996) Correlation Between Gingivain/Gingipain and Bacterial Dipeptidyl Peptidase Activity in Gingival Crevicular Fluid and Periodontal Attachment Loss in Chronic Periodontitis Patients. A 2-Year Longitudinal Study. Journal of Periodontology 67, 703-716.
- 10 Eley B.M. and Cox S.W. (2003) Proteolytic and hydrolytic enzymes from putative periodontal pathogens: characterization, molecular genetics, effects on host defences and tissues and detection in gingival crevice fluid. Periodontology 2000, 31, 105-124.
- 15 Jin L.J., Söder P.-Ö., Leung W.K., Corbet E.F., Samaranayake L.P., Söder B. and Davies W.I.R. (1999) Granulocyte elastase activity and PGE2 levels in gingival crevicular fluid in relation to the presence of subgingival periodontopathogens in subjects with untreated adult periodontitis. Journal of Clinical Periodontology 26, 531-540.
- 20 Johansson A., Sandström G., Claesson R., Hånström L., Kalfas S. (2000) Anaerobic neutrophil-dependent killing of Actinobacillus actinomycetemcomitans in relation to the bacterial leukotoxicity. European Journal of Oral Sciences 108, 136-149.
- 25 Lamster I.B., Smith Q.T., Celenti R.S., Singer R.E. and Grbic J.T. (1994) Development of a risk profile for periodontal disease: microbial and host response factors. Journal of Periodontology 65 (5 Suppl), 511-20.
- 30 Nakagawa, T., Sims, T., Fan, Q., Potempa, J., Travis, J., Houston, L. and Page, R.C. (2001) Functional characteristics of antibodies induced by Arg-gingipain (HRgpA) and Lys-gingipain (Kgp) from Porphyromonas gingivalis. Oral Microbiology and Immunology 16 (4), 202-211.
- 35

7:54 -01-30

Nisengard R.J., Mikulski L., McDuffie D. and Bronson P. (1992) Development of a rapid latex agglutination test for periodontal pathogens. Journal of Periodontology 63 (7), 611-617.

- 5 Oringer R.J. (2002) Modulation of the Host Response
in Periodontal Therapy. Journal of Periodontology 73,
460-470.

Socransky S.S., Smith C., Martin L., Paster B.J.,
Dewhirst F.E. and Levin A.E. (1994) "Checkerboard" DNA-
DNA hybridization. Biotechniques 17 (4), 788-92.

+46 31630263

Ink. I. Patent- och register

-01-30

24

Hjogdalen Kassin

CLAIMS

1. A test kit for detecting periodontal disease in a patient by analysing a sample from the oral cavity of the patient, wherein said kit at least comprises:

a first detection assay for detecting a first substance originating from bacteria, and

a second detection assay for detecting a second substance originating from the immune or inflammatory system of the patient.

2. A test kit according to claim 1, wherein said first detection assay comprises at least a first affinity ligand having a binding site for binding said first substance originating from bacteria, and

said second detection assay comprises at least a second affinity ligand having binding site for binding said second substance originating from the immune or inflammatory system of the patient.

3. A test kit according to claim 1 or 2, wherein said first substance is a bacterial virulence product.

4. A test kit according to claim 3, wherein said first substance is an enzyme.

5. A test kit according to claim 4, wherein said enzyme is a protease.

6. A test kit according to claim 5, wherein said protease is selected from the group consisting of arg-gingipain from *Porphyromonas gingivalis* and a 48 kDa protease from *Bacteroides forsythus*.

7. A test kit according to claim 3, wherein said first substance is a toxin.

+46 31630263

Ink. 1. Patent och register

04-01-30

25

Huvudlaxen Kassan

8. A test kit according to claim 7, wherein said toxin is a leukotoxin from *Actinobacillus actinomycetem-comitans*.

5 9. A test kit according to any of the preceding claims, wherein said second substance is a leukocyte product.

10 10. A test kit according to claim 9, wherein said leukocyte product is a natural serine protease.

11. A test kit according to claim 10, wherein said natural serine protease is a human neutrophil elastase.

15 12. A test kit according to any of the claims 1-8, wherein said second substance is a cytokine.

20 13. A test kit according to claim 12, wherein said cytokine is an interleukin.

14. A test kit according to claim 13, wherein said interleukin is chosen from among interleukin-1 β , interleukin-6 and interleukin-8.

25 15. A test kit according to claim 12, wherein said cytokine is an inflammatory mediator.

30 16. A test kit according to claim 15, wherein said inflammatory mediator is selected from the group consisting of tumour necrosis factor- α and prostaglandin E₂.

35 17. A test kit according to any of the claims 2 to 16, wherein said first affinity ligand is a first antibody exhibiting selective binding of said first substance and said second affinity ligand is a second antibody exhibiting selective binding of said second substance.

2004-01-30

Huvudfaxen Kassan

27

25. A method according to claim 24, wherein said first substance is a bacterial virulence product.

5 26. A method according to claims 25, wherein said first substance is an enzyme.

27. A method according to claim 26, wherein said enzyme is a protease.

10 28. A method according to claim 27, wherein said protease is selected from the group consisting of arg-gingipain from Porphyromonas gingivalis and a 48 kDa protease from Bacteroides forsythus.

15 29. A method according to claim 25, wherein said first substance is a toxin.

20 30. A method according to claim 29, wherein said toxin is a leukotoxin from Actinobacillus actinomycetemcomitans.

31. A method according to any of the claims 24-30, wherein said second substance is a leukocyte product.

25 32. A method according to claim 30, wherein said leukocyte product is a natural serine protease.

30 33. A method according to claim 32, wherein said natural serine protease is a human neutrophil elastase.

34. A method according to any of the claims 24-30, wherein said second substance is a cytokine.

35 35. A method according to claim 36, wherein said cytokine is an interleukin.

032

+46 31630263

Ink. t. Patent- och reg.verket

70614 -01- 3 0

Huvudfäken Kattor

1/1




Fig. 1

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
84

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☒ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:**

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.